




Edition: BP 2025 (Ph. Eur. 11.6 update)

## Glucosamine Sulfate Sodium Chloride



### General Notices

(Ph. Eur. monograph 2447)



$\text{C}_{12}\text{H}_{28}\text{Cl}_2\text{N}_2\text{Na}_2\text{O}_{14}\text{S}$  573.3 216447-62-0

Ph Eur

## DEFINITION

Bis(2-amino-2-deoxy-D-glucopyranose) sulfate bis(sodium chloride).

Substance prepared from glucosamine hydrochloride isolated from natural sources or produced by fermentation, and sodium sulfate.

### Content

98.0 per cent to 102.0 per cent (dried substance).

## PRODUCTION

The animals from which glucosamine sulfate sodium chloride is derived must fulfil the requirements for the health of animals suitable for human consumption.

## CHARACTERS

### Appearance

White or almost white, crystalline powder.

### Solubility

Freely soluble in water, sparingly soluble in methanol, practically insoluble in acetone.

## IDENTIFICATION

A. Infrared absorption spectrophotometry ([2.2.24](#)).

Comparison [glucosamine sulfate sodium chloride CRS](#).

B. It gives reaction (a) of chlorides ([2.3.1](#)).

C. 1 mL of solution S (see Tests) gives reaction (a) of sodium ([2.3.1](#)).

D. It gives reaction (a) of sulfates ([2.3.1](#)).

E. Specific optical rotation (see Tests).

## TESTS

### Solution S

Dissolve 2.50 g in [carbon dioxide-free water R](#) and dilute to 25.0 mL with the same solvent.

### Appearance of solution

The solution is clear ([2.2.1](#)) and colourless ([2.2.2, Method II](#)).

Dilute 5.0 mL of solution S to 25.0 mL with [water R](#).

### pH ([2.2.3](#))

3.0 to 5.0 for solution S.

### Specific optical rotation ([2.2.7](#))

+ 50.0 to + 55.0 (dried substance), determined on solution S.

Examine 3 h after preparation of solution S.

### Related substances

Liquid chromatography ([2.2.29](#)).

**Test solution** To 0.40 g of the substance to be examined add 80 mL of the mobile phase and sonicate for 10 min. Cool to room temperature and dilute to 100.0 mL with the mobile phase.

**Reference solution (a)** Dissolve 25.0 mg of [2-methylpyrazine CRS](#) in the mobile phase and dilute to 10.0 mL with the mobile phase. Dilute 1.0 mL of the solution to 10.0 mL with the mobile phase. Dilute 1.0 mL of this solution to 100.0 mL with the mobile phase.

**Reference solution (b)** Dissolve the contents of a vial of [glucosamine for system suitability CRS](#) (containing impurities B and C) in 1 mL of the mobile phase.

**Column:**

— size:  $l = 0.15$  m,  $\varnothing = 4.6$  mm;

— stationary phase: [base-deactivated end-capped octadecylsilyl silica gel for chromatography R](#) (3  $\mu$ m);

— temperature: 30 °C.

**Mobile phase** Dissolve 0.5 g of [sodium heptanesulfonate R](#) in [water for chromatography R](#), add 0.5 mL of [phosphoric acid R](#) and 4 mL of a 56 g/L solution of [potassium hydroxide R](#) and dilute to 1000 mL with [water for chromatography R](#); to 1000 mL of this solution add 50 mL of [acetonitrile R1](#).

**Flow rate** 1.0 mL/min.

**Detection** Spectrophotometer at 195 nm.

**Injection** 20 µL.

**Run time** Twice the retention time of 2-methylpyrazine.

**Retention time** 2-methylpyrazine = about 9 min.

**System suitability** Reference solution (b):

— **resolution**: minimum 1.5 between the peaks due to impurities B and C.

**Limits:**

— **unspecified impurities**: for each impurity, not more than 0.5 times the area of the principal peak in the chromatogram obtained with reference solution (a) (0.05 per cent);

— **total**: not more than twice the area of the principal peak in the chromatogram obtained with reference solution (a) (0.2 per cent);

— **disregard limit**: 0.3 times the area of the principal peak in the chromatogram obtained with reference solution (a) (0.03 per cent).

### **Loss on drying (2.2.32)**

Maximum 0.5 per cent, determined on 1.000 g by drying in an oven at 105 °C for 2 h.

### **Sulfated ash (2.4.14)**

23.5 per cent to 26.0 per cent, determined on 1.0 g.

Ignite first at 600 ± 50 °C for 2 h. Do not repeat the moistening with [sulfuric acid R](#) between any re-ignition.

### **Microbial contamination**

TAMC: acceptance criterion 10<sup>3</sup> CFU/g ([2.6.12](#)).

TYMC: acceptance criterion 10<sup>2</sup> CFU/g ([2.6.12](#)).

Absence of [Escherichia coli](#) ([2.6.13](#)).


## **ASSAY**

Dissolve 0.250 g in 50 mL of [water R](#) and add 1.0 mL of [0.1 M hydrochloric acid](#). Titrate with [0.1 M sodium hydroxide](#), determining the end-point potentiometrically ([2.2.20](#)). Read the volume added between the 2 points of inflexion.


1 mL of [0.1 M sodium hydroxide](#) is equivalent to 28.67 mg of C<sub>12</sub>H<sub>28</sub>Cl<sub>2</sub>N<sub>2</sub>Na<sub>2</sub>O<sub>14</sub>S.

## **IMPURITIES**


*Other detectable impurities (the following substances would, if present at a sufficient level, be detected by one or other of the tests in the monograph. They are limited by the general acceptance criterion for other/unspecified impurities and/or by the general monograph [Substances for pharmaceutical use \(2034\)](#). It is therefore not necessary to identify these impurities for demonstration of compliance. See also [5.10. Control of impurities in substances for pharmaceutical use](#))* A, B, C, E.




A. 2-(acetylamino)-2-deoxy-D-glucopyranose (*N*-acetylglucosamine),



B. (1*R*,1'*R*,2*S*,2'*S*,3*R*,3'*R*)-1,1'-pyrazine-2,5-diylbis (butane-1,2,3,4-tetrol) (fructosazine),



C. (1*R*,2*S*,3*R*)-1-[5-[(2*S*,3*R*)-2,3,4-trihydroxybutyl] pyrazin-2-yl]butane-1,2,3,4-tetrol (deoxyfructosazine),



E. 5-(hydroxymethyl)furan-2-carbaldehyde (5-hydroxymethylfurfural).

---

Ph Eur