



Edition: BP 2025 (Ph. Eur. 11.6 update)

Heavy Magnesium Carbonate



[General Notices](#)

(Ph. Eur. monograph 0043)

Action and use

Antacid; osmotic laxative.

Preparations

[Calcium Carbonate and Heavy Magnesium Carbonate Chewable Tablets](#)

[Compound Magnesium Trisilicate Oral Powder](#)

Ph Eur

DEFINITION

Hydrated basic magnesium carbonate.

Content

40.0 per cent to 45.0 per cent, calculated as MgO (M_r 40.30).

CHARACTERS

Appearance

White or almost white powder.

Solubility

Practically insoluble in water. It dissolves in dilute acids with effervescence.

IDENTIFICATION

- A. Untapped bulk density ([2.9.34](#)): minimum 0.25 g/mL.
- B. It gives the reaction of carbonates ([2.3.1](#)).

C. Dissolve about 15 mg in 2 mL of [dilute nitric acid R](#) and neutralise with [dilute sodium hydroxide solution R](#). The solution gives the reaction of magnesium ([2.3.1](#)).

TESTS

Solution S

Dissolve 5.0 g in 100 mL of [dilute acetic acid R](#). When the effervescence has ceased, boil for 2 min, allow to cool and dilute to 100 mL with [dilute acetic acid R](#). Filter, if necessary, through a previously ignited and tared porcelain or silica filter crucible of suitable porosity to give a clear filtrate.

Appearance of solution

Solution S is not more intensely coloured than reference solution B₄ ([2.2.2, Method II](#)).

Soluble substances

Maximum 1.0 per cent.

Mix 2.00 g with 100 mL of [water R](#) and boil for 5 min. Filter whilst hot through a sintered-glass filter (40) ([2.1.2](#)), allow to cool and dilute to 100 mL with [water R](#). Evaporate 50 mL of the filtrate to dryness and dry at 100-105 °C. The residue weighs not more than 10 mg.

Substances insoluble in acetic acid

Maximum 0.05 per cent.

Any residue obtained during the preparation of solution S, washed, dried, and ignited at 600 ± 50 °C, weighs not more than 2.5 mg.

Chlorides ([2.4.4](#))

Maximum 700 ppm.

Dilute 1.5 mL of solution S to 15 mL with [water R](#).

Sulfates ([2.4.13](#))

Maximum 0.6 per cent.

Dilute 0.5 mL of solution S to 15 mL with [distilled water R](#).

Calcium ([2.4.3](#))

Maximum 0.75 per cent.

Dilute 2.6 mL of solution S to 150 mL with [distilled water R](#). 15 mL of the solution complies with the test.

Iron ([2.4.9](#))

Maximum 400 ppm.

Dissolve 0.1 g in 3 mL of [dilute hydrochloric acid R](#) and dilute to 10 mL with [water R](#). Dilute 2.5 mL of the solution to 10 mL with [water R](#).

ASSAY

Dissolve 0.150 g in a mixture of 2 mL of [dilute hydrochloric acid R](#) and 20 mL of [water R](#). Carry out the complexometric titration of magnesium ([2.5.11](#)).

1 mL of [0.1 M sodium edetate](#) is equivalent to 4.030 mg of MgO.

FUNCTIONALITY-RELATED CHARACTERISTICS

This section provides information on characteristics that are recognised as being relevant control parameters for one or more functions of the substance when used as an excipient (see chapter [5.15](#)). Some of the characteristics described in the Functionality-related characteristics section may also be present in the mandatory part of the monograph since they also represent mandatory quality criteria. In such cases, a cross-reference to the tests described in the mandatory part is included in the Functionality-related characteristics section. Control of the characteristics can contribute to the quality of a medicinal product by improving the consistency of the manufacturing process and the performance of the medicinal product during use. Where control methods are cited, they are recognised as being suitable for the purpose, but other methods can also be used. Wherever results for a particular characteristic are reported, the control method must be indicated.

The following characteristics may be relevant for heavy magnesium carbonate used as filler in tablets.

Particle-size distribution ([2.9.31](#) or [2.9.38](#))

Bulk density of powders ([2.9.34](#))

Ph Eur