



Edition: BP 2025 (Ph. Eur. 11.6 update)

Paracetamol and Caffeine Soluble Tablets

[General Notices](#)

Soluble Paracetamol and Caffeine Tablets

Action and use

Analgesic; antipyretic; central nervous system stimulant.

DEFINITION

Paracetamol and Caffeine Soluble Tablets contain Paracetamol and Caffeine in either a suitable soluble basis or a suitable soluble, effervescent basis.

The tablets comply with the requirements stated under Tablets and with the following requirements.

Content of paracetamol, $C_8H_9NO_2$

95.0 to 105.0% of the stated amount.

Content of caffeine, $C_8H_{10}N_4O_2$

95.0 to 105.0% of the stated amount.

IDENTIFICATION

A. Shake a quantity of the powdered tablets containing 0.5 g of Paracetamol with 20 mL of [acetone](#), filter and evaporate the filtrate to dryness. The [infrared absorption spectrum](#) of the residue, [Appendix II A](#), is concordant with the reference spectrum of paracetamol ([RS 258](#)).

B. Carry out the method for [thin-layer chromatography](#), [Appendix III A](#), using the following solutions.

- (1) Mix with the aid of ultrasound a quantity of powdered tablets containing 65 mg of Caffeine in 10 mL of [methanol](#), and filter (a 1.2- μ m Whatman GF/C is suitable).
- (2) 0.65% w/v of [caffeine BPCRS](#) in [methanol](#).
- (3) 0.65% w/v of [caffeine BPCRS](#) and 5% w/v of [paracetamol BPCRS](#) in [methanol](#).

CHROMATOGRAPHIC CONDITIONS

- (a) Use as the coating [silica gel F₂₅₄](#) (Merck silica gel 60 F₂₅₄ plates are suitable).
- (b) Use the mobile phase as described below.
- (c) Apply 1 μ L of each solution.
- (d) Develop the plate to 15 cm.
- (e) After removal of the plate, allow it to dry in air and examine under [ultraviolet light \(254 nm\)](#).

MOBILE PHASE

5 volumes of [acetic acid](#), 5 volumes of [ethanol](#), 5 volumes of [water](#) and 50 volumes of [ethyl acetate](#).

SYSTEM SUITABILITY

The test is not valid unless the chromatogram obtained with solution (3) shows two clearly separated spots.

CONFIRMATION

The spot corresponding to caffeine in the chromatogram obtained with solution (1) corresponds in position to the principal spot in the chromatogram obtained with solution (2).

C. In the Assay for caffeine, the chromatogram obtained with solution (1) shows a peak with the same retention time as the principal peak in the chromatogram obtained with solution (2).

TESTS

Related substances

Carry out the method for [liquid chromatography, Appendix III D](#), using the following solutions prepared in solution A.

Solution A: 0.23% w/v of [sodium chloride](#) in a mixture of 30 volumes of mobile phase B and 70 volumes of mobile phase A.

- (1) Mix with the aid of ultrasound a quantity of the powdered tablets containing 0.5 g of Paracetamol with 50 mL and filter (Chromafil RC 45/25 is suitable).
- (2) Dilute 1 volume of solution (1) to 100 volumes.
- (3) 0.0001% w/v of [4-aminophenol](#) (paracetamol impurity K).
- (4) 0.00001% w/v of [4'-chloroacetanilide](#) (paracetamol impurity J).
- (5) Dilute 1 volume of solution (2) to 10 volumes.

CHROMATOGRAPHIC CONDITIONS

- (a) Use a stainless steel column (15 cm × 4.6 mm) packed with [end-capped octadecylsilyl silica gel for chromatography](#) (2.6 µm) (Kinetex C18 100A is suitable).
- (b) Use gradient elution and the mobile phase as described below.
- (c) Use a flow rate of 0.8 mL per minute.
- (d) Use a column temperature of 35°.
- (e) Use detection wavelengths of 212 nm and 246 nm.
- (f) Inject 20 µL of each solution.

MOBILE PHASE

Mobile phase A 5 mM [sodium octanesulfonate](#), adjusted to pH 2.2 with [orthophosphoric acid](#).

Mobile phase B [methanol R1](#).

Time (Minutes)	Mobile phase A (% v/v)	Mobile phase B (% v/v)	Comment
0-2.5	80→70	20→30	linear gradient
2.5-20	70	30	isocratic
20-30	70→20	30→80	linear gradient
30-32	20→80	80→20	linear gradient
32-37	80	20	re-equilibration

When the chromatograms are recorded under the prescribed conditions, the relative retentions with reference to paracetamol (retention time about 3 minutes) are: caffeine impurity B, about 0.6; caffeine impurity F, about 1.2; caffeine impurity A, about 1.3; caffeine, about 1.6; paracetamol impurity K, about 2.3; caffeine impurity E, about 2.5; paracetamol impurity J, about 6.1.

SYSTEM SUITABILITY

The test is not valid unless:

in the chromatogram obtained with solution (3) at 212 nm, the [signal-to-noise ratio](#) of the peak due to paracetamol impurity K is at least 10.

in the chromatogram obtained with solution (4) at 246 nm, the [signal-to-noise ratio](#) of the peak due to paracetamol impurity J is at least 10.

LIMITS

For paracetamol impurity J at 246 nm

In the chromatogram obtained with solution (1):

the area of any peak corresponding to paracetamol impurity J is not greater than the area of the principal peak in the chromatogram obtained with solution (4) (10 ppm).

For all other impurities at 212 nm

Identify any peaks due to caffeine impurity B and E, and multiply the peak areas by a correction factor of 2.9 and 3.3, respectively.

In the chromatogram obtained with solution (1):

the area of any peak corresponding to paracetamol impurity K is not greater than the area of the corresponding peak in the chromatogram obtained with solution (3) (100 ppm);

the area of any other [secondary peak](#) is not greater than the area of the peak due to paracetamol in the chromatogram obtained with solution (5) (0.1%);

The total impurity content is not greater than 0.5%.

Disregard any peak, excluding paracetamol impurities J and K, with an area less than half the area of the peak due to paracetamol in the chromatogram obtained with solution (5) (0.05%).

ASSAY

For paracetamol

Weigh and powder 20 tablets. Carry out the method for [liquid chromatography](#), [Appendix III D](#), using the following solutions protected from light.

- (1) Shake a quantity of the powdered tablets containing 0.5 g of Paracetamol with 100 mL of the mobile phase, dilute to 200 mL with the same solvent, filter through a glass-fibre filter (Whatman GF/C is suitable) and dilute 5 mL of the filtrate to 250 mL with the mobile phase.
- (2) 0.005% w/v of [paracetamol BPCRS](#) in the mobile phase.

CHROMATOGRAPHIC CONDITIONS

- (a) Use a stainless steel column (10 cm × 4.6 mm) packed with [octadecylsilyl silica gel for chromatography](#) (5 µm) (Nucleosil C18 is suitable).
- (b) Use isocratic elution and the mobile phase described below.
- (c) Use a flow rate of 1.5 mL per minute.
- (d) Use an ambient column temperature.
- (e) Use a detection wavelength of 243 nm.
- (f) Inject 20 µL of each solution.

MOBILE PHASE

0.01M [sodium pentanesulfonate](#) in a mixture of 22 volumes of [methanol](#) and 78 volumes of [water](#), the pH of the solution being adjusted to 2.8 using [2M hydrochloric acid](#).

DETERMINATION OF CONTENT

Calculate the content of C₈H₉NO₂ in the tablets from the chromatograms obtained and using the declared content of C₈H₉NO₂ in [paracetamol BPCRS](#).

For caffeine

Weigh and powder 20 tablets. Carry out the method for [liquid chromatography](#), [Appendix III D](#), using the following solutions protected from light.

- (1) Shake a quantity of the powdered tablets containing 30 mg of Caffeine with 100 mL of the mobile phase, filter through a glass-fibre filter (Whatman GF/C is suitable) and dilute 5 mL of the filtrate to 50 mL with the mobile phase.
- (2) 0.003% w/v of [caffeine BPCRS](#) in the mobile phase.

CHROMATOGRAPHIC CONDITIONS

The chromatographic conditions described under the Assay for paracetamol may be used with a detection wavelength of 220 nm.

DETERMINATION OF CONTENT

Calculate the content of $C_8H_{10}N_4O_2$ in the tablets from the chromatograms obtained and using the declared content of $C_8H_{10}N_4O_2$ in [caffeine BPCRS](#).

LABELLING

The label states (1) that the tablets should be dissolved in water immediately before use; (2) where applicable, that the tablets are effervescent; (3) the quantities of Paracetamol and Caffeine.

IMPURITIES

The impurities limited by the requirements of this monograph include impurities J and K listed under [Paracetamol](#), and impurities A, B, E and F listed under [Caffeine](#).